


Cambodia


Total population (2016): 15,827 241 ▶ Population aged 15 years and older (15+): 69% ▶ Population in urban areas: 21% ▶ Income group (World Bank): Lower-middle income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	2010*		2016*	
Recorded	2.0		3.5	
Unrecorded	3.0		3.2	
Total**	5.0		6.7	
Total males / females	8.8	1.5	11.9	2.0
WHO Western Pacific Region	7.0		7.3	

* Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

	Litres
Males (15+)	27.4
Females (15+)	10.1
Both sexes (15+)	21.7

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	20.6	47.7	14.0	51.0
Females	3.2	16.3	1.9	18.3
Both sexes	11.5	37.3	8.1	42.1

* Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	30.3	59.6	45.6
Former drinkers* (15+)	26.5	20.8	23.5
Abstainers (15+), past 12 months	56.8	80.4	69.1

* Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	105.0	61.3	66.5	38.0	3 399
Road traffic injuries, males / females	42.4	19.0	26.1	15.3	687
Cancer, males / females	211.3	145.1	5.4	1.6	466

*Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016

LEAST < 1 2 3 4 **5** > MOST

* Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	8.7	5.0
Females	1.8	0.7
Both sexes	5.1	2.7
WHO Western Pacific Region	4.7	2.3

* 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Yes (2013/—) / Yes
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	No / No / No
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	No / No / No
Restrictions for on-/off-premise sales of alcoholic beverages (any):	
Hours, days / places, density	No, No / No, No
Specific events / intoxicated persons / petrol stations	No / No / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.05 / 0.05 / 0.05
Legally binding regulations on alcohol advertising / product placement (any)	Yes / No
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / No
Legally required health warning labels on alcohol advertisements / containers (any)	No / No
National government support for community action (any)	Yes
National monitoring system(s) (any)	No